

AD-A039 262

OHIO STATE UNIV RESEARCH FOUNDATION COLUMBUS
ROLE OF CARBON DIOXIDE IN INERT GAS NARCOSIS.(U)
APR 77 H S WEISS
OSURF-3916-F

F/G 6/19

UNCLASSIFIED

N00014-67-A-0232-0025
NL

1 OF 1
AD
A039262



END

DATE
FILMED
5-77



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

ADAU39262

RF Project 3916

Report No.

12
NW

FINAL

REPORT

By

THE OHIO STATE UNIVERSITY
RESEARCH FOUNDATION

1314 KINNEAR RD.
COLUMBUS, OHIO 43212

To DEPARTMENT OF THE NAVY
Office of Naval Research
Arlington, Virginia 22217

N00014-67-A-0232-0025

On ROLE OF CARBON DIOXIDE IN INERT GAS NARCOSIS

For the period May 1, 1974 - December 31, 1976

Submitted by Dr. Harold S. Weiss

Department of Physiology

Date April, 1977

D D C

RECEIVED
MAY 11 1977
D

DISTRIBUTION STATEMENT A

Approved for public release;

Reproduction in whole or in part
is permitted for any purpose of
the United States Government.

FILE COPY

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER (9) Final 4 rept. 1 May 74-31 Dec 76	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) ROLE OF CARBON DIOXIDE IN INERT GAS NARCOSIS	5. TYPE OF REPORT & PERIOD COVERED Final 5/1/74 - 12/31/76	
7. AUTHOR (10) Harold S. Weiss	6. PERFORMING ORG. REPORT NUMBER 3916-F Final	
9. PERFORMING ORGANIZATION NAME AND ADDRESS The Ohio State University Research Foundation, 1314 Kinnear Road Columbus, Ohio 43212	8. CONTRACT OR GRANT NUMBER(s) N00014-67-A-0232-0025	
11. CONTROLLING OFFICE NAME AND ADDRESS Department of the Navy Office of Naval Research Arlington, Virginia 22217	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) (12) 6p.	12. REPORT DATE (11) Apr 77	
	13. NUMBER OF PAGES 4	
	15. SECURITY CLASS. (of this report) Unclassified	
	15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited. (14) OSURF-3916-F		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) carbon dioxide, oxygen, high pressure narcosis, avian, unidirectional flow, eliminates dead space, PCO_2 and PO_2 and $P(\text{inert})$ maintained, visually evoked response, depression of VER appears to be a reflection of narcosis induced by high nitrogen pressure, body temperature depressed VER		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The role of carbon dioxide and oxygen in high pressure narcosis was studied by exposing animals to hyperbaric conditions while maintaining them normoxic and normocapnic. Chickens were the experimental animals, used since the avian respiratory anatomy lends itself to a surgical modification that allows a unidirectional flow of gas through the pulmonary system and eliminates dead space. Heated, humidified gas entered the lung via a tracheal cannula and exited from the posterior air sacs via cannulae through the body wall. If the mass flow of gas through the lungs is much greater than the normal flow, then the bird's own respiratory		

DD FORM 1473
1 JAN 73

EDITION OF 1 NOV 65 IS OBSOLETE

Unclassified

(con't.)

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

20.

gas exchange will not significantly affect the composition of the gas. Therefore the lung PCO_2 and PO_2 and $P(\text{inert})$ can be maintained at any level chosen for the inflow gas. The unanaesthetized restrained birds were exposed to pressures up to 6 atmospheres absolute (ATA) in a hyperbaric chamber.

→ Narcosis was estimated from changes in the visually evoked response (VER). The VER was recorded from the optic tectum by means of implanted bipolar fine wire electrodes. The VER was initiated by an external strobe light. A depression of the amplitude of the VER is considered to reflect narcosis.

All animals showed a strong depression in VER amplitude on exposure to normoxic, normocapnic gas at 6 ATA when the inert component was nitrogen; the onset of the depression was a smooth function of depth. No depression of VER was seen when the inert diluent was helium. Increasing lung PCO_2 to 70 mmHg had no effect on VER. Therefore depression of VER appears to be a reflection of narcosis induced by high nitrogen pressure and not solely hypoxia, hypercapnia or pressure per se. Relatively small changes in the body temperature (1.5 C) depressed VER and may explain some of the depression of VER seen by others with helium. Resistance to the high flow of gas through the bird was usually 7.5 - 10 cm H_2O at a flow rate of 6 l/min ambient temperature and pressure; this resistance remains nearly constant over the 1 - 6 ATA pressure range investigated.

ACCESSION for	
WIB	White Section <input checked="" type="checkbox"/>
OCG	Buff Section <input type="checkbox"/>
UNRECORDED	<input type="checkbox"/>
JUSTIFICATION	
BY	
DISPOSITION/AVAILABILITY CODES	
AVAIL. NO./IN SPECIAL	
A	

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

The Role of Carbon Dioxide in Inert Gas Narcosis

Narcosis induced by breathing compressed air at elevated pressures has been known since the 1800s but the mechanism of action is still debated. The most widely held theory considers narcosis to be a direct effect of the increased quantity of nitrogen (N_2) dissolved in tissues under high pressure. The dissolved N_2 is believed to interfere with neural function by changing permeability and/or electrical polarization of cells and cellular membranes. This theory appears to fit in well with the Meyer-Overton view of anesthesia.

A competing hypothesis suggests that the neural disturbances are brought about by hypercapnia and hypoxia. It is presumed that hyperbaria increases gas density and viscosity to the point where respiratory flow is inhibited, leading to retention of carbon dioxide (CO_2) and a deficiency of oxygen (O_2). Acceptance of this theory appeared to be declining, in part because gases like xenon are narcotic at sea-level pressures and in part because N_2 narcosis was not alleviated in man when ventilation was assisted by pump. However, it has been restimulated by an analysis which suggests that rather than simple inhibition of flow, subsegmental bronchioles may collapse on expiration when gas of increased density flows through them at velocities associated with normal ventilatory rates. The collapsed passages would effectively reduce the diffusion area and lead to hypercapnia and hypoxia. This revised theory can also explain the xenon effects at sea-level pressure, and the failure of narcosis to be relieved by a respiratory pump in the circuit.

Experimentally, it has been difficult to substantiate or refute the hypoxia/hypercapnia concept, in part because of problems associated with measuring oxygen and carbon dioxide tensions (PO_2 , PCO_2) in arterial blood while at elevated pressures. Our experimental approach was somewhat different; namely, to use the unique characteristics of the avian lung to establish any desired concentration of gases at the blood-gas exchange surface. This is possible in Aves because pulmonary gas flow can be made unidirectional through the interconnecting network of air capillaries and air sacs which replace the mammalian alveoli and diaphragm. Pulmonary PCO_2 , PO_2 , and PN_2 can thus be controlled independently at any desired pressure and flow and with no dead space.

In this study the role of carbon dioxide and oxygen in high-pressure narcosis was examined by exposing unanesthetized, mildly restrained chickens to hyperbaric conditions while maintaining them normoxic and normocapnic. The chickens were surgically modified to allow the unidirectional flow of heated, humidified gas to enter the lung via a tracheal cannula and to exit from the posterior air sacs via cannulae through the body wall. The mass flow of gas through the lungs was 10 times the normal flow, effectively preventing the bird's own respiratory gas exchange from significantly altering the composition of the pulmonary gas. PCO_2 , PO_2 , and P_{inert} in the lung was therefore maintained at the level chosen for the inflow gas.

The unanesthetized, restrained birds were exposed to pressures up to six atmospheres absolute (ATA) in a hyperbaric chamber. Narcosis was estimated from changes in the visually evoked response (VER). The VER was recorded from the optic tectum of the brain by means of implanted bipolar fine-wire electrodes. The VER was initiated by an external strobe light. A depression of the amplitude of the VER was considered to reflect narcosis.

All animals showed a strong depression in VER amplitude on exposure to normoxic, normocapnic gas at 6 ATA when the inert component was N_2 ; the onset of the depression was a smooth function of depth. No depression of VER was seen when the inert diluent was helium (He). In birds at 6 ATA with N_2 , increasing lung P_{O_2} to the equivalent of 60% O_2 at sea level and decreasing lung P_{CO_2} below 4% sea-level equivalent did not bring about any recovery of the depressed VER. Therefore depression of VER appears to be a reflection of narcosis induced by high nitrogen pressure and not of hypoxia, hypercapnia, or pressure per se.

Minimal effects of CO_2 were confirmed when increasing lung P_{CO_2} to 70 mmHg or decreasing it to near zero mmHg had no demonstrable effect on the VER of the bird at sea level. The avian brain however, seems very sensitive to hypoxia, since decreasing lung P_{O_2} only to the 120-130 mmHg range depressed the VER. Relatively small changes in the body temperature ($1.5^\circ C$) also depressed the VER. In view of the high heat conductivity of He, this may explain some of the reports of narcotic effects with helium. Resistance to the high flow of gas through the bird was usually 7.5 - 10 cm H_2O at a flow rate of 6 l/min at ambient temperature and pressure; this resistance remains nearly constant over the 1-6 ATA pressure range investigated.

Lack of a satisfactory animal preparation appears to have inhibited experimental work on inert gas narcosis. The awake, unidirectionally ventilated chicken prepared for recording the VER appears to be an animal model well suited for such studies. It may serve to carry out comparisons among inert gases, on various combinations of O_2 (particularly hypoxia) and CO_2 with the inert gases, and also for evaluating a range of pharmacological and physical factors (e.g., temperature) believed important in narcosis.

PUBLICATIONS

1. Torley, L. and H.S. Weiss, "Effect of Age and Magnesium Ions on Oxygen Toxicity in the Neonate Chicken Gallus Domesticus." Undersea Biomed. Res. 2:223-227, 1975.
2. Torley, L. and H.S. Weiss, "Visual Evoked Response in Gallus Domesticus exposed to high pressure with known lung gas compositions." (Abst.) Fed. Proc. 34(3):462, 1975.

3. Torley, L. and H.S. Weiss, "Use of an Avian Preparation to Study the Role of CO₂ in Nitrogen Narcosis. (Abst.) 8th Symposium on Underwater Physiology; San Diego, California, July, 1975.
4. Torley, L. "The Role of Hypercapnia and Hypoxia in High Pressure Narcosis." Ph.D. Dissertation, The Ohio State University, 1975.